Burock MA, Buckner RL, Dale AM (1998) Understanding differential responses in event related fMRI through linear simulation. In: Proc ISMRM 6th Annu Meeting, Sydney, p. 245

Butts K, Riederer SJ, Ehman RL, Thompson RM, Jack CR (1994) Interleaved echo planar imaging on a standard MRI system. Magn Reson Med 31:67–72

Buxton RB, Luh WM, Wong EC, Frank LR, Bandettini PA (1998a) Diffusion-weighting attenuates the BOLD signal change but not the post-stimulus undershoot. In: Proc ISMRM 6th Annu Meeting, Sydney, p. 7

Buxton RB, Wong EC, Frank LR (1998b) Dynamics of blood flow and oxygenation changes during brain activation: the balloon model. Magn Reson Med 39:855–864

Cao Y, Vikingstad EM, Huttenlocher PR, Levin DN (1994) Functional magnetic resonance studies of the reorganization of the human hand sensorimotor area after unilateral brain injury in the perinatal period. Proc Natl Acad Sci USA 91:9612–9616

Clark VP, Maisog JM, Haxby JV (1998) fMRI study of face perception and memory using random stimulus sequences. J Neurophys 79:3257–3265

Cohen JD, Perlstein WM, Braver TS, Nystrom LE, Noll DC, Jonides J et al. (1997) Temporal dynamics of brain activation during a working memory task. Nature 386:604–607

Cohen MS (1997) Parametric analysis of fMRI data using linear systems methods. Neuroimage 6:93–103

Cohen MS, Weisskoff RM (1991) Ultra-fast imaging. Magn Reson Imaging 9:1–37

Courtney SM, Ungerleider LG, Keil K, Haxby JV (1997) Transient and sustained activity in a distributed neural system for human working memory. Nature 386:608–611

Dale AM, Buckner RL (1997) Selective averaging of rapidly presented individual trials using fMRI. Hum Brain Mapp 5:329–340

Davis TL, Weisskoff RM, Kwong KK, Savoy R, Rosen BR (1994) Susceptibility contrast undershoot is not matched by inflow contrast undershoot. In: Proc SMR 2nd Annu Meeting, San Francisco

Davis TL, Kwong KK, Weisskoff RM, Rosen BR (1998) Calibrated functional MRI: mpping the dynamics of oxidative metabolism. Proc Natl Acad Sci USA 95:1834–1839

Detre JA, Leigh JS, WIlliams DS, Koretsky AP (1992) Perfusion imaging. Magn Reson Med 23:37–45

DeYoe EA, Bandettini P, Neitz J, Miller D, Winans P (1994) Functional magnetic resonance imaging (FMRI) of the human brain. J Neurosci Methods 54:171–187

Duyn JH, Moonen CTW, vanYperen GH, de Boer RW, Luyten PR (1994) Inflow versus deoxyhemoglobin effects in BOLD functional MRI using gradient-echoes at 1.5 T. NMR Biomed 7:83–88

Edelman RR, Sievert B, Wielopolski P, Pearlman J, Warach S (1994) Noninvasive mapping of cerebral perfusion by using EPISTAR MR angiography. J Magn Reson Imaging 4(P):68 (abstract)

Frahm J, Bruhn H, Merboldt K-D, Hanicke W, Math D (1992) Dynamic MR imaging of human brain oxygenation during rest and photic stimulation. J Magn Reson Imaging 2:501–505

Frahm J, Merboldt K-D, Hanicke W (1993) Functional MRI of human brain activation at high spatial resolution. Magn Reson Med 29:139–144

Frahm J, Krüger G, Merboldt K-D, Kleinschmidt A (1996) Dynamic uncoupling and recoupling of perfusion and oxidative metabolism during focal activaiton in man. Magn Reson Med 35:143–148

Fransson P, Krüger G, Merboldt K-D, Frahm J (1998) Temporal characteristics of oxygenation-sensitive responses to visual activation in humans. Magn Reson Med 39:912–919

Friston KJ, Jezzard P, Turner R (1994) Analysis of functional MRI time-series. Hum Brain Mapp 2:69–78

Friston KJ, Fletcher P, Josephs O, Holmes A, Rugg MD, Turner R (1998a) Event-related fMRI: characterizing differential responses. Neuroimage 7:30–40

Friston KJ, Josephs O, Rees G, Turner R (1998b) Nonlinear event-related responses in fMRI. Magn Reson Med 39:41–52

Glover GH, Lee AT (1995) Motion artifacts in fMRI: comparison of 2DFT with PR and spiral scan methods. Magn Reson Med 33:624–635

Grinvald A, Frostig RD, Siegel RM, Bratfeld E (1991) High-resolution optical imaging of functional brain architecture in the awake monkey. Proc Natl Acad Sci USA 88:11559–11563

Haacke EM, Lai S, Yablonski DA, Lin W (1995) In vivo validation of the BOLD mechanism: a review of signal changes in gradient-echo functional MRI in the presence of flow. Int J Imag Syst Technol 6:153–163

Haacke EM, Lai S, Reichenbach JR, Kuppusamy K, Hoogenraad FGC, Takeichi H et al. (1997) In vivo measurement of blood oxygen saturation using magnetic resonance imaging: a direct validation of the blood oxygen level-dependent concept in functional brain imaging. HIum Brain Mapp 5:341–346

Hennig J, Janz C, Speck O, Ernst T (1995) Functional spectroscopy of brain activation following a single light pulse: examinations of the mechanism of the fast initial response. Int J Imag Syst Technol 6:203–208

Hickok G, Love T, Swinney D, Wong EC, Buxton RB (1997) Functional MR imaging during auditory word perception: a single-trial presentation paradigm. Brain Language 58: 197–201

Hu X, Kim S-G (1994) Reduction of signal fluctuations in functional MRI using navigator echoes. Magn Reson Med 31:495–503

Hu X, Le TH, Ugurbil K (1997) Evaluation of the early response in fMRI in individual subjects using short stimulus duration. Magn Reson Med 37:877–884

Janz C, Schmitt C, Fischer H, Speck O, Hennig J (1998) Variability of the hemodynamic response to different stimuli in single event paradigms. In: Proc ISMRM 6th Annu Meeting, Sydney

Jesmanowicz A, Bandettini PA, Hyde JS (1998) Single-shot half k-space high-resolution gradient-recalled EPI for fMRI at 3 tesla. Magn Reson Med 40:754-762

Josephs O, Turner R, Friston K (1997) Event-related fMRI. Hum Brain Mapp 5:243–248

Karni A, Meyer G, Jezzard P, Adams MM, Turner R, Ungerleider LG (1995) Functional MRI evidence for adult motor cortex plasticity during motor skill learning. Nature 377:155–158

Kennan RP, Zhong J, Gore JC (1994) Intravascular susceptibility contrast mechanisms in tissues. Magn Reson Med 31:9–21

Kim S-G (1995) Quantification of relative cerebral blood flow change by flow-sensitive alternating inversion recovery

(FAIR) technique: application to functional mapping. Magn Reson Med 34:293–301
Kim S-G, Ugurbil K (1997) Comparison of blood oxygenation and cerebral blood flow effects in fMRI: estimation of relative oxygen consumption change. Magn Reson Med 38:59–65
Kim S-G, Ashe J, Georgopoulos AP, Merkle H, Ellermann JM, Menon RS (1993) Functional imaging of human motor cortex at high magnetic field. J Neurophysiol 69:297–302
Kim S-G, Richter W, Ugurbil K (1997) Limitations of temporal resolution in fucntional MRI. Magn Reson Med 37:631–636
Konishi S, Yoneyama R, Itagaki H, Uchida I, Nakajima K, Kato H et al. (1996) Transient brain activity used in magnetic resonance imaging to detect functional areas. Neuroreport 8: 19–23
Kwong KK, Belliveau JW, Chesler DA, Goldberg IE, Weisskoff RM, Poncelet BP et al. (1992) Dynamic magnetic resonance imaging of human brain activity during primary sensory stimulation. Proc Natl Acad Sci USA 89:5675–5679
Kwong KK, Chesler DA, Weisskoff RM, Rosen BR (1994) Perfusion MR imaging. In: Proc SMR 2nd Annu Meeting, San Francisco
Le TH, Hu X (1996) Retrospective estimation and correction of physiological artifacts in fMRI by direct extraction of physiological activity form MR data. Magn Reson Med 35:290–298
Lee AT, Glover GH, Meyer CH (1995) Discrimination of large venous vessels in time-course spiral blood-oxygen-level-dependent magnetic-resonance functional neuroimaging. Magn Reson Med 33:745–754
Li S-J, Jesmanowicz A, Bodurka J, Hyde J (1998) Current-induced magnetic resonance phase imaging. In: Proc ISMRM 6th Annu Meeting, Sydney, p.1495
Luknowsky DC, Gati JS, Menon RS (1998a) Mental chronometry using single trials and EPI at 4 T. In: Proc ISMRM 6th Annu Meeting, Sydney, p.167
Mandeville JB, Marota JJA, Kosofky BE, Keltner JR, Weissleder R, Rosen BR et al. (1998) Dynamic functional imaging of relative cerebral blood volume during rat forepaw stimulation. Magn Reson Med 39:615–624
McCarthy G, Luby M, Gore J, Goldman-Rakic P (1997) Infrequent events transiently activate human prefrontal and parietal cortex as measured by functional MRI. J Neurophysiol 77:1630–1634
McKinnon GC (1993) Ultrafast interleaved gradient-echo-planar imaging on a standard scanner. Magn Reson Med 30:609–616
Menon RS, Ogawa S, Tank DW, Ugurbil K (1993) 4 Tesla gradient recalled echo characteristics of photic stimulation-induced signal changes in the human primary visual cortex. Magn Reson Med 30:380–386
Menon RS, Ogawa S, Strupp JP, Anderson P, Ugurbil K (1995a) BOLD based functional MRI at 4 tesla includes a capillary bed contribution: echo-planar imaging correlates with previous optical imaging using intrinsic signals. Magn Reson Med 33:453–459
Menon RS, Ogawa S, Ugurbil K (1995b) High-temporal-resolution studies of the human primary visual cortex at 4 T: teasing out the oxygenation contribution in FMRI. Int J Imag Syst Technol 6:209–215
Moonen CTW, van Zijl PCM, Frank JA, LeBihan D, Becker ED (1990) Functional magnetic resonance imaging in medicine and physiology. Science 250:53–61
Noll DC (1995) Methodologic considerations for spiral k-space functional MRI. Int J Imag Syst Technol 6:175–183
Ogawa S, Lee TM (1992) Functional brain imaging with physiologically sensitive image signals. J Magn Reson Imaging 2(P)-WIP (Suppl):S22 (abstract)
Ogawa S, Lee TM, Kay AR, Tank DW (1990) Brain magnetic resonance imaging with contrast dependent on blood oxygenation. Proc Natl Acad Sci USA 87:9868–9872
Ogawa S, Tank DW, Menon R, Ellermann JM, Kim S-G, Merkle H et al. (1992) Intrinsic signal changes accompanying sensory stimulation: functional brain mapping with magnetic resonance imaging. Proc Natl Acad Sci USA 89:5951–5955
Ogawa S, Menon RS, Tank DW, Kim S-G, Merkle H, Ellerman JM et al. (1993) Functional brain mapping by blood oxygenation level-dependent contrast magnetic resonance imaging: a comparison of signal characteristics with a biophysical model. Biophys J 64:803–812
Richter W, Andersen PM, Georgopolous AP, Kim S-G (1997a) Sequential activity in human motor areas during a delayed cued finger movement task studied by time-resolved fMRI. Neuroreport 8:1257–1261
Richter W, Ugurbil K, Georgopouloos A, Kim S-G (1997b) Time resolved fMRI of mental rotation. Neuroreport 8:3697–3702
Rosen BR, Belliveau JW, Chien D (1989) Perfusion imaging by nuclear magnetic resonance. Magn Reson Q 5:263–281
Rosen BR, Buckner RL, Dale AM (1998) Event-related functional MRI: past, present, future. Proc Natl Acad Sci USA 95:773–780
Saad ZS, Ropella KM, Carman GJ, DeYoe EA (1996) Temporal phase variation of FMR signals in vasculature versus parenchyma. In: Proc ISMRM 4th Annu Meeting, New York
Sakai K, Hikosaka O, Miyauchi S, Takino R, Sasaki Y, Putz B (1998) Transition of brain activation from frontal to parietal areas in visuomotor sequence learning. J Neurosci 18:1827–1840
Savoy RL, O'Craven KM, Weisskoff RM, Davis TL, Baker J, Rosen B (1994) Exploring the temporal bourdaries of fMRI: measuring responses to very brief visual stimuli. In: Book of Abstracts, Soc Neurosci 24th Annu Meeting, Miami
Savoy RL, Bandettini PA, Weisskoff RM, Kwong KK, Davis TL, Baker JR et al. (1995) Pushing the temporal resolution of fMRI: studies of very brief visual stimuli, onset variablity and asynchrony, and stimulus-correlated changes in noise. In: Proc SMR 3rd Annu Meeting, Nice, p.450
Schacter DL, Buckner RL (1998) Priming and the brain. Neuron 20:185–195
Schacter DL, Buckner RL, Koutstaal W, Dale AM, Rosen BR (1997) Late onset of anterior prefrontal activity during true and false recognition: an event related fMRI study. Neuroimage 6:259–269
Schmitt F, Stehling MK, Turner R (1998) Echo-planar imaging: theory, technique, and application. Springer, Berlin Heidelberg New York
Song AW, Wong EC, Tan SG, Hyde JS (1996) Diffusion weighted fMRI at 1.5 T. Magn Reson Med 35:155–158
Stehling MK, Turner R, Mansfield P (1991) Echo-planar imaging: magnetic resonance imaging in a fraction of a second. Science 254:43–50

Turner R, LeBihan D, Moonen CTW, Despres D, Frank J (1991) Echo-planar time course MRI of cat brain oxygenation changes. Magn Reson Med 22:159–166

van Zijl PCM, Eleff SM, Ulatowski JA, Oja JME, Ulug AM, Traystman RJ, Kauppinen RA (1998) Quantitative assessment of blood flow, blood volume and blood oxygenation effects in functional magnetic resonance imagning. Nature Med 4:159–167

Vasquez A, Noll D (1998) Nonlinear aspects of the BOLD response in functional MRI. Neuroimage 7:108–118

Williams DS, Detre JA, Leigh JS, Koretsky AS (1992) Magnetic resonance imaging of perfuusion using spin-inversion of arterial water. Proc Natl Acad Sci USA 89:212–216

Wong EC, Buxton RB, Frank LR (1997) Implementation of quantitative perfusion imaging techniques for functional brain mapping using pulsed arterial spin labeling. NMR Biomed 10:237–249

Wong EC, Buxton RB, Frank LR (1998) Quantitative imaging of perfusion using a single subtraction (QUIPSS and QUIPSSII). Magn Reson Med 39:702–708

Ye FQ, Smith AM, Yang Y, Duyn J, Mattay VS, Ruttimann UE et al. (1997) Quantitation of regional cerebral blood flow increases during motor activation: a steady-state arterial spin tagging study. Neuroimage 6:104–112

Zarahn E, Aguirre G, D'Esposito M (1997) A trial-based experimental design for fMRI. Neuroimage 6:122–138